

Minimal standards for describing new species belonging to the families *Campylobacteraceae* and *Helicobacteraceae*: *Campylobacter*, *Arcobacter*, *Helicobacter* and *Wolinella* spp.

Stephen L. W. On,^{1,*} William G. Miller,² Kurt Houf,^{3,4} James G. Fox⁵ and Peter Vandamme⁴

Abstract

Ongoing changes in taxonomic methods, and in the rapid development of the taxonomic structure of species assigned to the Epsilonproteobacteria have lead the International Committee of Systematic Bacteriology Subcommittee on the Taxonomy of *Campylobacter* and Related Bacteria to discuss significant updates to previous minimal standards for describing new species of *Campylobacteraceae* and *Helicobacteraceae*. This paper is the result of these discussions and proposes minimum requirements for the description of new species belonging to the families *Campylobacteraceae* and *Helicobacteraceae*, thus including species in *Campylobacter*, *Arcobacter*, *Helicobacter*, and *Wolinella*. The core underlying principle remains the use of appropriate phenotypic and genotypic methods to characterise strains sufficiently so as to effectively and unambiguously determine their taxonomic position in these families, and provide adequate means by which the new taxon can be distinguished from extant species and subspecies. This polyphasic taxonomic approach demands the use of appropriate reference data for comparison to ensure the novelty of proposed new taxa, and the recommended study of at least five strains to enable species diversity to be assessed. Methodological approaches for phenotypic and genotypic (including whole-genome sequence comparisons) characterisation are recommended.

INTRODUCTION

The class Epsilonproteobacteria is a phylogenetically-distinct lineage within the Proteobacteria [1] and currently contains 16 genera (Tables S1 and S2, available with the online version of this article). The genus *Campylobacter* was the first of these established [2], and the realisation that certain species were important human and animal pathogens prompted many further studies investigating the wider ecology and distribution of similar organisms. Bacteria found in cases of human gastritis and initially classified as *Campylobacter* spp. were later reclassified into a separate but related genus, *Helicobacter* [3]. Improvements in isolation, detection and taxonomic characterization methods, together with continued interest in the significance and distribution of such bacteria have resulted in the present status of the Epsilonproteobacteria – a highly diverse group of organisms containing over 100 taxa.

Within the class, phylogenetic subgroups can be identified. Of these, most taxa in the families *Campylobacteraceae*

(namely *Campylobacter*, *Arcobacter* and *Sulfurospirillum*) and *Helicobacteraceae* (namely *Helicobacter* and *Wolinella*) appear more closely related to each other than to free-living Epsilonproteobacteria such as the Nautiliaceae [4]. Their known or potential significance as pathogens, and advances in cultivation, make *Campylobacter* and *Helicobacter* the most populous genera, with many taxa sharing the same ecological niche. Their close phenotypic similarity to each other is illustrated by many *Arcobacter*, *Sulfurospirillum* and *Helicobacter* spp. having originally been described as ‘*Campylobacter*’ species (reviewed in [5]).

In every year since 1988, at least one novel species belonging to at least one of these genera has been described. On occasion, descriptions have been controversial [6–9], and on several occasions the ICSP Subcommittee on the Taxonomy of *Campylobacter* and Related Bacteria have remarked on the sometimes questionable quality of particular species descriptions that were observed [10–12]. Furthermore, taxonomic methods have undergone a ‘sea change’ with whole-

Author affiliations: ¹Department of Wine, Food and Molecular Biosciences, Lincoln University, PO Box 85084, Lincoln, New Zealand; ²U.S. Department of Agriculture, Produce Safety and Microbiology Research Unit, Agricultural Research Service, Albany, CA, USA; ³Department of Veterinary Public Health, Faculty of Sciences, Ghent University, K. L. Ledeganckstraat 35, B-9000 Ghent, Belgium; ⁴Department of Biochemistry and Microbiology, Laboratory of Microbiology, Faculty of Sciences, Ghent University, K. L. Ledeganckstraat 35, B-9000 Ghent, Belgium; ⁵Department of Comparative Medicine, Massachusetts Institute of Technology, 77, Massachusetts Avenue, Cambridge, MA 02139, USA.

***Correspondence:** Stephen L. W. On, stephen.on@lincoln.ac.nz

Keywords: minimal standards; species descriptions; polyphasic taxonomy; *Campylobacteraceae*; *Helicobacteraceae*; *Arcobacter*; *Wolinella*; Epsilonproteobacteria.

Abbreviations: ANI, average nucleotide identity; GBDP, Genome Blast Distance Phylogeny; TMAO, trimethylamine N-oxide. Two supplementary tables are available with the online version of this article.

genome sequencing offering a new and more detailed perspective on organismal relationships.

This paper accounts for these substantive changes and expands and updates previously-published minimal standards for describing new species of *Campylobacteraceae* [13] and *Helicobacter* [14]. Table S1 lists the current, validly-described species names in the genera covered by this proposal with their type strains and accompanying 16S rRNA gene sequence accession number. Members of the genus *Sulfurospirillum* are exclusively free-living and, from an ecological perspective, have more in common with, for example, members of the *Nautilaceae* (cf. Table S2), for which minimal standards will be proposed in due course. Intriguingly, phylogenetic analyses based on both 16S rRNA gene sequences (Fig. 1), and concatenated protein sequences derived from a subset of 289 conserved single-copy-number genes reveal a close relationship between *Sulfurospirillum* and *Campylobacter* [15], despite these taxa inhabiting radically different ecological niches. Such observations highlight the complexities of the Epsilonproteobacteria group as a whole, and support the need for a polyphasic taxonomic approach for accurate classification.

GENERAL FEATURES OF *CAMPYLOBACTERACEAE*

Gram-negative, curved, spiral or occasionally straight rod-shaped cells, 0.2–5 µm long. Cells may become more spherical after time or with exposure to environmental stress. Non-spore forming. Most species are motile by use of one to two unsheathed polar flagella, but a few species are aflagellate. Optimum growth temperatures range from 25 °C (notably *Arcobacter* spp.) to 42 °C (some *Campylobacter* spp.). All *Campylobacter* spp. can be cultured at 37 °C under appropriate conditions. The optimal growth atmosphere for *Campylobacteraceae* is microaerobic – traditionally, such atmospheres are described as comprising 3–8 % O₂, with commonly available Gaspak systems providing the balance with 10 % CO₂ and the remainder as N₂. However, some species require H₂ for growth and studies have demonstrated all *Campylobacter* spp. can grow in an atmosphere of 3 % O₂, 7 % H₂, 10 % CO₂ and 80 % N₂. Most *Arcobacter* spp. demonstrate the ability to grow under aerobic conditions (*A. anaerophilus* is an exception as an obligate anaerobe) [16], but all arcobacters to date can grow at 20–30 °C. Most species produce oxidase (undetected in *C. gracilis* and in some strains of *C. showae* [17]) and, in conventional laboratory testing for oxidation and/or fermentation of glucose, (e.g. [18]), do not demonstrate the ability to ferment or oxidise carbohydrates. *Campylobacter* and some *Arcobacter* spp. may be associated with the oral environment and/or the enteric or reproductive tracts of host animals; some *Arcobacter* spp. are free-living [4, 5, 16].

GENERAL FEATURES OF *HELICOBACTERACEAE*

Gram-negative, spiral, helical, curved, or fusiform rods of width 0.3–0.6 µm and length 1–5 µm. Cells may become more spherical after time or with exposure to environmental stress. Non-spore forming. Most species are motile by means of single or multiple flagella. In most species the flagella are sheathed. Optimum temperatures of growth range from 37 to 42 °C and all known species can be cultured at 37 °C. They are usually microaerophilic and best cultured in atmospheres containing H₂. Oxidase producing with most strains producing catalase. Gastric *Helicobacter* spp. described to date predominantly produce copious amounts of urease. Using common laboratory methods (e.g. [18]), *Helicobacteraceae* do not demonstrate the ability to ferment or oxidise carbohydrates. *Helicobacter* and *Wolinella* species described thus far have been associated with gastric, enteric, reproductive and/or hepatic environments of a variety of host animal species [1, 3, 7, 14, 19, 20]; none are known to be free-living.

General comments

The description of new species belonging to *Campylobacteraceae* or *Helicobacteraceae* should be based on characteristics necessary for assigning the new taxon to the genus, and on characteristics serving to differentiate the new taxon from existing taxa of the genus. In practice, this will require a polyphasic taxonomic approach utilizing both genotypic and phenotypic methods, especially since there is no single phenotypic characteristic that readily enables the assignment of strains to the genera *Campylobacter*, *Arcobacter*, *Helicobacter* or *Wolinella*. Furthermore, species such as *C. coli* and *H. pullorum* may be found in the same environments (presently chickens and humans) and share many phenotypic characters, rendering them difficult to discriminate using conventional phenotyping [5]. Thus, a phylogenetic assignment based on, at least, comparative analysis of 16S rRNA gene sequences is mandatory to appropriately assign strains to genus level [5, 21]. Corresponding sequences from type strains of relevant, validly-named Epsilonproteobacterial taxa should be included in sequence comparisons to appropriately designate phylogeny and thus genus placement. Additional gene sequence comparisons, such as *atpA* [22], *rpoB* [23] or *groEL* (Hsp60) [24], may provide useful information for strain classification and a finer resolution of taxonomic identity.

The description should ideally be based on not fewer than five isolates from different sources, or five distinct genotypes (i.e. representing distinguishable, individual strains) from the same or similar sources. The most useful taxonomic descriptions involve studies where species heterogeneity can be adequately assessed.

For critical comparisons with other species, controls consisting of type or reference strains of the appropriate taxa must be tested. For all phenotypic test procedures, the inoculum size, composition of the gaseous atmosphere,




Fig. 1. Phylogenetic relationships between 16S rRNA genes of type strains of the taxa encompassed in this study inferred by neighbor-joining tree using the Kimura 2-parameter distance estimation method, with bootstrapping based on 500 replicates.

temperature and period of incubation, and composition of the basal growth medium should be stated. The use of standardized, well-described tests and methods is recommended to facilitate comparison [25–29]. For descriptions at the

species level, the use of advanced phenotypic methods, such as and including MALDI-TOF MS analysis, should be regarded as supportive data but not the primary means by which species are delineated at the phenotypic level. Present

studies, although limited to relatively few taxa, indicate this approach has promise [30–33] with some caveats [34].

Putative new species of uncultured organisms for which molecular sequence data (such as 16S rRNA sequence) is available may qualify for assignment to the provisional taxonomic status *Candidatus* [35] in accordance with the proposals of Murray and Stackebrandt [36], as exemplified by the initial description of *C. hominis* prior to its culture [37].

Cell morphology

The reaction of cells in the Gram-staining procedure must be stated. The shape, size, and spiral wavelength (where appropriate) of bacterial cells should be reported. The tendency to undergo transformation to coccoid forms on exposure to air or in older cultures should be noted and the time taken for cells to change their appearance provided. The number and arrangement of flagella should be determined by electron microscopy, as well as the presence or absence of flagellar sheaths and periplasmic fibers.

Motility

Cells should be observed by microscopic examination of wet mounts or hanging drop preparations of young cultures in buffered saline or broth.

Growth conditions

Factors affecting growth should be tested under conditions that are near optimal unless stated otherwise.

- (1) Colony morphology. The size, shape, and colour of colonies should be described for optimal growth conditions on solid media. The type of nutrient agar base medium used (brain-heart infusion [BHI], Mueller-Hinton etc.) and the manufacturer from which it was obtained should be stated. The presence of swarming on solid media should be noted. When cultured on blood-containing agar media, the percentage and species of blood (e.g. horse, sheep, cattle) should be stated and any haemolytic activity described.
- (2) Temperature range. The time of incubation and ability to grow in specified broth or agar media from standardized inocula at various temperatures should be reported. The following temperatures should be used: 25, 37 and 42 °C.
- (3) Gaseous requirements. The ability of the strains to grow under aerobic, microaerobic and anaerobic conditions should be reported. The oxygen and hydrogen content must be specified for microaerobic conditions. The means by which such conditions were produced (e.g., gas replacement method, commercial gas-generating sachet [state manufacturer], incubator) must be given.

Biochemical properties

Results for the following tests are required: (i) oxidase activity, by use of any conventional method [28]; (ii) catalase activity, with percentage of reagent solution and time of observation given; (iii) nitrate reduction, preferably by the

plate method of Cook [38]; (iv) indoxyl acetate hydrolysis, preferably using a disc method [28] with percentage of reagent solution, volume of impregnation and time of observation given; (v) urease activity, using a rapid method [39]; (vi) alkaline phosphatase activity [40], with time of observation given; (vii) hippurate hydrolysis [41]; and (viii) selenite reduction [28]. Growth on media containing the following compounds using standardized methods [26, 27] should be determined: 2.0 % NaCl, 1 % glycine and 0.04 % triphenyl tetrazolium chloride (TTC); the ability of strains to reduce the latter should also be recorded when growth is observed.

Other tests

Test results for the following are desirable: γ -glutamyl transpeptidase [25]; growth on media containing 3.5 % NaCl, 0.032 % methyl orange, and 0.1 % sodium fluoride [26, 27, 29]; and anaerobic growth on 0.1 % trimethylamine N-oxide (TMAO) [27].

Resistance to antimicrobial agents

Susceptibility to nalidixic acid (30 μ g) and cephalothin (30 μ g), should be determined either by disc-diffusion or plate MIC tests. For diffusion assays, the absence of a clear zone of inhibition should be recorded as resistance; for susceptible strains, the inhibition zone sizes should be stated. The type of base medium used should be stated. Mueller-Hinton agar with added 10 % horse – or sheep – blood is recommended. Standardised procedures should be employed [26, 27, 42]. Brucella Agar and other media with bisulfite have been shown to inhibit the growth of *H. pylori* and therefore should be avoided [43].

Summary Tables for recommended phenotypic tests for *Arcobacter*, *Campylobacter*, *Helicobacter* and *Wolinella* spp. are provided (Tables 1, 2 and 3).

Phylogenetic analyses

The essentially complete (greater than 1450 bases) 16S rRNA sequence must be determined for the type strain, and ideally at least four additional independent isolates of the putative new species or subspecies. Specific methods for sequencing 16S rRNA from *Helicobacter* species, including intervening sequences (IVS), have been described [44]. Intervening sequences in the 16S rRNA gene should be fully sequenced and the 16S rRNA gene sequence, including any IVS, deposited in nucleic acid databases as a single sequence. Deposition of a complete genome sequence accomplishes this goal de facto. The phylogenetic position of representative strains of the putative new taxon must be determined by comparative sequence analysis of the 16S rRNA macromolecule [21]. Phylogenetic tree construction should demonstrate that the novel sequence clusters with those of all validly-named taxa of the appropriate genus within the *Campylobacteraceae* or *Helicobacteraceae*. A full description of alignment and treeing methods, including software, algorithms, treatment of gap penalties, and treatment of IVS sequences should be included. An exemplar tree is presented in Fig. 1.

Table 1. Summary phenotypic data for extant *Arcobacter* spp.

Data are derived from original species descriptions [16, 50, 74–88] (cf. Table S1) and/or On et al. [17].

	1	2	3	4	5	6	7	8	9	10
Growth temperature range (°C)	18–37	18–37	18–37	18–37	18–37	18–37	18–25,[37, 42]	18–37,42*	18–30	18–37, 42†±
Atmospheric requirements	ANO2	O2, mO2, ANO2†	O2, mO2, ANO2	O2, mO2, ANO2	[O2], mO2, ANO2*	O2, mO2, mO2	O2, mO2, [ANO2]	O2, mO2, ANO2†	O2, mO2	O2, mO2, ANO2±
Oxidase	–	+	+	+	+	+	+	+	+	+
Catalase	–	+	+	V	V	+	+	±	–	+
Nitrate reduction	+	+	–	+	–	+	V	+	–	+
Indoxyl acetate hydrolysis	+	+	+	+	+	+	+	+	+	+
Urease	–	–	–	–	–	–	–	+	+	V
Alkaline phosphatase	U	U	U	–	–	U	–	U	U	U
Hippuricase	U	U	U	–	–	U	–	U	U	U
Selenite reduction	U	U	U	–	–	U	–	U	U	U
Growth on: 2 % NaCl	+	+	+	M	–	+	M	+	+	+
1 % glycine	+	–	–	–	–	–	–	–	–	–
0.04 % TTC	–	–	–	+	V	+	M	–	–	–
TTC reduction	–	–	–	+	–	U	M	–	–	–
Resistance to: Nalidixic acid (30 mg)	U	U	U	F	V	U	–	U	U	U
Cephalothin (30 mg)	U	U	U	+	+	U	+	U	U	U
Desirable features:										
g-glutamyl transpeptidase	U	U	U	U	U	U	U	U	U	U
Growth on:										
3.5 % NaCl	+	U	+	V	–	+	–	U	+	U
0.032 % methyl orange	U	U	U	+	+	U	+	U	U	U
0.1 % sodium fluoride	U	U	U	+	V	U	+	U	U	U
Anaerobic growth on 0.1 % TMAO	U	U	U	+	±	U	F	U	U	U

	11	12	13	14	15	16	17	18	19	20	21
Growth temperature range (°C)	18–37	25–37	10–40	18–37,42*±	18–37,42†±	18–25	18–37,[42]	18–30	18–30	18–30	18–37
Atmospheric requirements	O2, mO2, ANO2	O2, mO2	O2, mO2	O2, mO2, ANO2±	O2, mO2, ANO2†	O2, mO2, ANO2	O2, mO2, ANO2	O2, mO2	O2, mO2	O2, mO2	O2, mO2
Oxidase	+	+	+	+	+	+	+	+	+	+	1
Catalase	–	+	–	+	+	+	+	+	+	+	1
Nitrate reduction	+	+	+	+	+	+	+	+	+	–	1
Indoxyl acetate hydrolysis	+	+	+	–	–	+	+	+	+	+	1
Urease	–	U	–	–	–	+	–	–	–	–	1
Alkaline phosphatase	–	–	–	U	+	–	–	U	–	U	U
Hippuricase	–	–	–	U	U	–	–	U	–	–	U
Selenite reduction	–	U	U	U	U	V	F	U	–	+	U

Table 1. cont.

	11	12	13	14	15	16	17	18	19	20	21
Growth on: 2 % NaCl	+	U	+	+	+	+	+	–	M	V	1
1 % glycine	–	+	+	–	+	–	–	–	+	–	–
0.04 % TTC	–	U	–	–	–	–	+	–	V	–	–
TTC reduction	–	+	–	–	+	–	+	–	V	–	–
Resistance to: Nalidixic acid (30 mg)	–	–	U	U	U	–	–	U	V	U	U
Cephalothin (30 mg)	–	U	U	U	U	–	+	U	+	+	U
Desirable features:											
g-glutamyl transpeptidase	U	–	U	U	U	U	U	U	U	U	U
Growth on:											
3.5 % NaCl	+	U	+	+	+	+	+	–	V	V	+
0.032 % methyl orange	–	U	U	U	U	+	+	U	+	U	U
0.1 % sodium fluoride	–	U	U	U	U	+	+	U	+	U	U
Anaerobic growth on 0.1 % TMAO	U	U	U	U	U	+	–	U	V	U	U

1, *A. anaerophilus*; 2, *A. aquimarinus*; 3, *A. bivalviorum*; 4, *A. butzleri*; 5, *A. cibarius*; 6, *A. cloacae*; 7, *A. cryaerophilus*; 8, *A. defluvi*; 9, *A. ebronensis*; 10, *A. ellisi*; 11, *A. halophilus*; 12, *A. lanthieri*; 13, *A. marinus*; 14, *A. molluscorum*; 15, *A. mytili*; 16, *A. nitrofigilis*; 17, *A. skirrowii*; 18, *A. suis*; 19, *A. therius*; 20, *A. trophiarum*; 21, *A. venerupis*. + = all strains examined give a positive result. – = all strains examined give a negative result. O2, aerobic conditions; ANO2, anaerobic conditions. () = variable (31–58 %) number of strains grow at this temperature; [] = few (11–25 %) strains grow in these conditions; V, 33–67 % strains positive; ±, weak activity; + = 02 only; t, at 30 °C only; F, 22–25 % strains positive; M, 84–95 % strains positive; U, unknown at this time.

The 16S rRNA gene is the most widely used phylogenetic marker but there are known problems with the accurate phylogenetic placement of certain species within the *Campylobacteraceae* or *Helicobacteraceae* [7, 45–47]. The use of additional phylogenetic markers such as *atpA* [22], *rpoB* [23] or *groEL* [24], may provide useful data to support the phylogenetic position of the new taxon. Multi-Locus Sequence Analysis [48] or rMLST [49] may provide a more robust description of the taxonomic position of a given strain. Taxonomic studies of *Campylobacteraceae* or *Helicobacteraceae* species using MLSA and rMLST are as yet uncommon, although MLSA has been applied to *Arcobacter* species [32, 50]. Ultimately, these approaches can be expected to be replaced by whole genome sequence based phylogenomic analyses (e.g. [15, 51, 52]). As with 16S rRNA gene sequence comparisons, representatives of all extant validly-published named taxa in the genus in which the proposed novel organism is placed must be included.

Genomic analyses

A complete genome sequence of the proposed type strain of a novel species or subspecies is a definitive means to determine the guanosine-plus-cytosine (% G+C) content of the DNA; however, classical methods (melting temperature or enzymatic) may also be used [53]. Reference DNA such as *Escherichia coli* ATCC 11775^T (G+C, 51 mol%) or *H. pylori* ATCC 700392 (G+C, 39 mol%) should be analysed at the same time and its estimated G+C content (moles percent) expressed relative to the reference DNA should be reported.

The current taxonomic definition of a species requires determination of the whole-genomic similarity, whereby at or around 70 % DNA–DNA relatedness indicates strains are sufficiently related to be assigned to the same species [54–56]. Methodological caveats notwithstanding (discussed below), novel species should demonstrate genomic relatedness to extant species at values discernibly lower than this level. The proposed type strain for any new taxon must be used in these comparisons.

Whole-genome relatedness can be tested in several ways. Classical DNA–DNA hybridization experiments have been used for many years (e.g. [57–60]). It is recognized that experiments on Epsilonproteobacteria can be difficult and that different methods can result in differing estimates of the degree of relatedness between strains [61, 62]. Numerical comparison of whole-cell protein profiles [5, 6, 60], and high-resolution amplified fragment length polymorphism (AFLP)-based fingerprints has also been shown to be effective in accurately determining genetic relationships between strains [63, 64]. It is essential that analyses are performed against a database of strain profiles of sufficient number and quality that represent all taxa with validly described species names, for results to be meaningful.

Where whole-genome sequences of proposed new species are available, *in silico* analyses that mimic conventional DNA–DNA hybridisations can be used in comparisons with type strain genomes of validly-published extant species,

Table 2. Summary phenotypic data for extant *Campylobacter* spp.

Data are derived from original species descriptions [6, 31, 60, 67–69, 87–120] (cf. Table S1) and/or On et al. [17].

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Growth temperature range (°C)	37–42	37–42	30–42	(30), 37, (42)	25–42	37, (42)	30, 37, (42)	(18–22), 25–37, (42)	25–37, (42)	<18–22>, (25), 37 mO ₂ , (ANO2)	(30), 37, (42) ANO ₂ , [HmO ₂]	37–42 mO ₂	[30], 37–42 mO ₂	37 ANO ₂ , [HmO ₂]	[18–25], 30–42 mO ₂	30–42 mO ₂ , ANO ₂
Atmospheric requirements	+	+	+	+	+	+	+	+	+	+	–	+	+	+	+	+
Oxidase	+	+	+	+	+	+	+	+	+	+	–	+	+	+	+	+
Catalase	+	+	+	+	+	+	–	+	+	+	+	+	–	–	+	+
Nitrate reduction	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Indoxyl acetate hydrolysis	+	+	+	+	+	+	+	–	–	–	+	+	+	–	–	–
Urease	–	–	–	–	+	–	–	–	–	–	–	–	–	–	–	–
Alkaline phosphatase	U	–	–	–	+	–	–	–	–	–	–	–	–	–	–	–
Hippuricase	+	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Selenite reduction	U	U	+	+	U	–	–	–	–	–	–	–	–	–	–	–
Growth on: 2% NaCl	U	–	–	–	+	–	–	–	–	–	–	–	–	–	–	–
1% glycine	+	+	+	+	+	–	+	+	+	+	+	+	+	+	+	+
0.04% TTC	U	U	+	–	–	–	–	–	–	–	–	–	–	–	–	–
TTC reduction	U	U	+	–	–	–	–	–	–	–	–	–	–	–	–	–
Resistance to: Nalidixic acid (30 mg)	U	U	+	–	–	–	–	–	–	–	–	–	–	–	–	–
Cephalothin (30 mg)	U	U	+	–	–	–	–	–	–	–	–	–	–	–	–	–
Desirable features:	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
g-glutamyl transpeptidase	–	+	U	U	U	–	U	U	U	U	U	U	U	U	U	U
Growth on:	U	–	–	–	U	–	–	–	–	–	–	–	–	–	–	–
3.5% NaCl	U	–	–	–	U	–	–	–	–	–	–	–	–	–	–	–
0.032% methyl orange	U	U	+	–	U	U	–	+	U	–	–	–	–	–	–	–
0.1% sodium fluoride	U	U	+	–	U	U	–	–	U	–	–	–	–	–	–	–
Anaerobic growth on 0.1% TMAO	U	U	–	–	U	U	+	–	U	–	+	–	–	+	–	–

	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Growth temperature range (°C)	18–37	37	37	(30), 37–42	37–42	37–42	30–42	30–42	37–42	37–42	37–42	37	<30>, 37, <42>	30–37, [42]	(30), 37, [42]	37–42	30–37, [42]	30–37, [42]	37–42
Atmospheric requirements	mO ₂ , ANO ₂ *	mO ₂	mO ₂	mO ₂	mO ₂ , ANO ₂ *	mO ₂	mO ₂	mO ₂ , ANO ₂	mO ₂ , ANO ₂	mO ₂	mO ₂ , ANO ₂	mO ₂ , ANO ₂	ANO ₂ , [HmO ₂]	ANO ₂ , [HmO ₂]	mO ₂ , ANO ₂	mO ₂ , ANO ₂	ANO ₂ , [HmO ₂]	ANO ₂ , [HmO ₂]	mO ₂ , ANO ₂ *
Oxidase	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Catalase	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Nitrate reduction	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Indoxyl acetate hydrolysis	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Urease	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–

Table 2. cont.

	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Alkaline phosphatase	U	U	-	-	+	U	-	M	-	U	U	U	-	-	-	-	-	-	-
Hippuricase	-	-	+	M	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Selenite reduction	U	+	-	M	U	U	V	F	U	U	U	U	-	-	V	-	+	-	+
Growth on: 2% NaCl	U	-	-	-	U	+	M	M	U	M	U	U	V	+	+	+	-	+	-
1 % glycine	+	+	F	M	-	+	+	V	+	+	-	-	+	V	+	M	+	+	-
0.04 % TTC	U	+	V	M	U	U	M	-	U	U	U	U	-	-	-	U	V	-	-
TTC reduction	U	+	V	M	U	U	M	-	F	U	U	U	-	-	-	U	V	-	-
Resistance to: Nalidixic acid (30 mg)	+	+	-	-	+	-	V	M	U	M	-	-	M	-	M	+	-	-	+
Cephalothin (30 mg)	-	+	-	M	+	+	+	F	U	F	-	-	-	-	-	-	F	-	+
Desirable features:																			
g-glutamyl transpeptidase	+	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U	U
Growth on:																			
3.5 % NaCl	U	-	-	-	U	-	-	-	U	U	U	U	-	-	V	-	-	+	-
0.032 % methyl orange	U	-	+	+	U	U	+	+	U	U	U	U	-	-	+	U	+	+	+
0.1 % sodium fluoride	U	-	F	+	U	U	+	-	U	U	U	U	-	+	M	+	-	+	+
Anaerobic growth on 0.1 % TMAO	U	V	-	-	U	U	+	+	U	U	U	U	+	V	M	+	-	+	U

1, *Campylobacter avium*; 2, 3, *Campylobacter coli*; 4, *Campylobacter concisus*; 5, *Campylobacter coraciensis*; 6, *Campylobacter cuniculorum*; 7, *Campylobacter curvus*; 8, *Campylobacter fetus* subsp. *fetus*; 9, *Campylobacter fetus* subsp. *testudinum*; 10, *Campylobacter fetus* subsp. *venerealis*; 11, *Campylobacter hepaticus*; 12, *Campylobacter gracilis*; 13, *Campylobacter helveticus*; 14, *Campylobacter hominis*; 15, *Campylobacter hyointestinalis* subsp. *hyointestinalis*; 16, *Campylobacter hyointestinalis* subsp. *lawsonii*; 17, *Campylobacter iguaniorum*; 18, *Campylobacter insulaenigrae*; 19, *Campylobacter jejuni* subsp. *jejuni*; 20, *Campylobacter jejuni* subsp. *jejuni*; 21, *Campylobacter lanienae*; 22, *Campylobacter lari* subsp. *concheus*; 23, *Campylobacter lari* subsp. *lari*; 24, *Campylobacter mucosalis*; 25, *Campylobacter ornithocola*; 26, *Campylobacter peloridis*; 27, *Campylobacter pinnipediorum* subsp. *pinnipediorum*; 28, *Campylobacter pinnipediorum* subsp. *caledonicus*; 29, *Campylobacter rectus*; 30, *Campylobacter showae*; 31, *Campylobacter sputorum*; 32, *Campylobacter subantarcticus*; 33, *Campylobacter upsaliensis*; 34, *Campylobacter ureolyticus*; 35, *Campylobacter volucris*. +=all strains examined give a positive result. -=all strains examined give a negative result <=>=20–30 % strains grow at this temperature; []=50–60 % strains grow at this temperature; ()=76–93 % grow at these conditions; 02, aerobic conditions; m02, microaerobic conditions; AN02, anaerobic conditions; Hm02=microaerobic atmosphere enhanced with H₂; F=7–27 % strains positive; V=29–57 % strains positive; M=70–95 % strains positive; *Weak growth.; †Biovar fecalis strains produce catalase; ‡Biovar paraureolyticus strains produce urease; §Urease-Positive Thermophilic *Campylobacter* (UPTC) variants.

Table 3. Summary phenotypic data for extant *Helicobacter* and *Wolinella* spp. Data are derived from original species descriptions [3, 19, 42, 44, 46, 57, 58, 62, 87, 121–154] (cf. Table S1) and/or On et al. [17].

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
Growth temperature range (°C)	(30), 37, <42>	37–42	37–42	37	37	37–42	37–42	37–42	<30>, 37–42	37–42	37–42	37–42	37–42	37	37	[30], 37, (42)	37	37
Atmospheric requirements	mO2	mO2	mO2	mO2, CO2, ANO2*	mO2	mO2	mO2	mO2	mO2	mO2	mO2, ANO2	mO2	mO2, ANO2	mO2	mO2	mO2	ANO2	mO2, ANO2*
Oxidase	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Catalase	+	+	+	+	+	+	+	+	–	+	+	M	+	+	+	M	F	+
Nitrate reduction	–	–	–	–	–	–	–	V	–	–	+	+	+	+	+	–	+	+
Indoxyl acetate hydrolysis	–	+	+	–	–	+	+	+	–	–	–	F	–	–	+	+	–	–
Urease	+	+	+	+	+	+	–	–	–	–	–	–	–	–	+	–	–	+
Alkaline phosphatase	+	–	–	+	U	+	–	–	+	–	U	F	+	+	+	V	–	–
Hippuricase	–	U	U	–	–	–	U	U	–	U	U	–	–	–	–	–	–	+
Selenite reduction	–	U	U	U	U	–	U	U	–	U	U	–	U	U	–	–	–	U
Growth on: 2% NaCl	–	U	–	–	–	–	U	–	–	U	–	–	–	–	–	–	–	–
1% glycine	–	+	–	–	+	–	+	–	–	U	–	–	+	–	–	–	–	–
0.04% TTC	–	U	U	U	+	–	U	–	V	U	U	+	U	U	–	+	+	U
TTC reduction	–	U	U	+	U	+	U	–	V	U	U	+	U	–	–	+	+	+
Resistance to: Nalidixic acid (30 mg)	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cephalothin (30 mg)	–	+	+	+	+	–	+	+	F	+	+	+	+	+	–	–	+	U
Desirable features:																		
g-glutamyl transpeptidase	U	–	+	+	U	+	–	–	U	+	–	U	–	–	U	U	U	+
Growth on:																		
3.5% NaCl	–	U	–	–	–	–	U	–	–	U	–	U	U	–	–	–	–	U
0.032% methyl orange	F	U	U	U	U	–	U	+	–	U	U	+	U	U	–	V	M	U
0.1% sodium fluoride	mO2	U	U	U	U	–	U	+	–	U	U	–	U	U	–	V	+	U
Anaerobic growth on 0.1% TMAO	–	U	U	U	U	–	U	+	–	U	U	–	U	U	–	–	+	U

	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Growth temperature range (°C)	37	30–42	37–42	37–42	37–42	37–42	37–42	37–42	37	37, (42)	37–42	37–42	(30), 37, <42>	37, [42]	37–42	37	37	37–42	37–42	37–42	37–42
Atmospheric requirements	mO2, ANO2	HmO2, ANO2	mO2	mO2	mO2	mO2	mO2	mO2, ANO2*	mO2	mO2, (ANO2)	mO2, ANO2*	mO2, ANO2	mO2, ANO2	mO2, ANO2	mO2	mO2, (ANO2)	mO2, ANO2*	mO2	mO2	mO2	ANO2
Oxidase	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Catalase	+	+	+	+	+	+	+	+	+	+	+	M	+	+	+	+	+	+	+	+	+
Nitrate reduction	+	+	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Indoxyl acetate hydrolysis	+	–	+	–	–	–	–	–	–	–	–	–	F	–	–	F	–	U	–	–	–
Urease	+	–	+	–	–	–	–	–	–	–	–	–	M	–	–	+	+	+	–	V	–
Alkaline phosphatase	U	+	–	–	–	–	–	–	–	–	–	–	+	–	–	V	+	F	–	–	–

Table 3. cont.

	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Hippuricase	–	–	U	U	U	U	U	–	–	–	–	–	–	–	U	–	–	–	–	–	–
Selenite reduction	U	U	U	U	U	U	U	U	–	+	U	+	–	U	U	–	U	U	U	U	+
Growth on: 2% NaCl	U	–	U	U	U	U	U	–	–	–	U	–	–	U	U	–	– ^a	–	– ^a	U	+
1% glycine	+	–	+	–	+	+	+	–	–	–	+	–	–	+	+	–	–	–	+	+	–
0.04% TTC	+	U	U	U	U	U	U	–	–	+	U	V	V	U	U	–	U	U	U	U	–
TTC reduction	U	–	U	U	U	U	U	–	–	+	U	V	V	U	U	–	+	U	U	–	–
Resistance to: Nalidixic acid (30 mg)	+	–	–	+	+	+	+	–	–	–	–	+	M	+	+	–	U	+	–	V	–
Cephalothin (30 mg)	+	+	+	+	+	+	+	+	–	+	+	–	F	+	+	V	U	+	+	+	–
Desirable features:																					
g-glutamyl transpeptidase	U	+	–	–	–	–	–	–	U	U	U	U	U	–	+	+	+	+	–	–	U
Growth on:																					
3.5% NaCl	U	U	U	U	U	U	U	–	–	–	–	–	–	U	U	–	– ^a	– ^a	– ^a	U	–
0.032% methyl orange	U	U	U	U	U	U	U	U	–	–	U	+	–	U	U	–	–	U	U	U	+
0.1% sodium fluoride	U	U	U	U	U	U	U	U	–	+	U	–	+	U	U	V	–	U	U	U	+
Anaerobic growth on 0.1% TMAO	+	U	U	U	U	U	U	U	–	–	U	–	F	U	U	V	–	U	U	U	+

1, *Helicobacter acinonychis*; 2, *Helicobacter anseris*; 3, *Helicobacter aurati*; 4, *Helicobacter baculiformis*; 5, *Helicobacter bilis*; 6, *Helicobacter bizzozeronii*; 7, *Helicobacter brantiae*; 8, *Helicobacter canadensis*; 9, *Helicobacter canis*; 10, *Helicobacter cetorum*; 11, *Helicobacter cholecystus*; 12, *Helicobacter cinaedi*; 13, *Helicobacter cynogastricus*; 14, *Helicobacter equorum*; 15, *Helicobacter fells*; 16, *Helicobacter fennelliae*; 17, *Helicobacter ganmani*; 18, *Helicobacter heilmannii*; 19, *Helicobacter hepaticus*; 20, *Helicobacter himalayensis*; 21, *Helicobacter jaachi*; 22, *Helicobacter japonicus*; 23, *Helicobacter macacae*; 24, *Helicobacter marmotae*; 25, *Helicobacter mastomyrinus*; 26, *Helicobacter mesocricetorum*; 27, *Helicobacter muridarum*; 28, *Helicobacter mustelae*; 29, *Helicobacter pametensis*; 30, *Helicobacter pullorum*; 31, *Helicobacter pylori*; 32, *Helicobacter rodentium*; 33, *Helicobacter saguini*; 34, *Helicobacter salomonis*; 35, *Helicobacter suis*; 36, *Helicobacter trogonum*; 37, *Helicobacter typhlonius*; 38, *Helicobacter valdiviensis*; 39, *Wolinella succinogenes*. +=all strains examined give a positive result. –=all strains examined give a negative result. O2, aerobic conditions; mO2, microaerobic conditions; ANO2, anaerobic conditions. <=>=18–27% strains positive; ()=33–67% strains positive; []=77–92% strains positive; F=6–29% strains positive; V=33–58% strains positive; M=83–93% strains positive; *, Weak growth; †, when API method used; I, intermediate resistance; a, assumed; no growth on 1.5% NaCl media.

Table 4. Exemplar results of whole-genome comparisons for well-characterized strains of *Campylobacter*, *Aerobacter* and *Helicobacter* using Average Nucleotide Identity (ANI) [65] and Genome Blast Distance Phylogeny (GBDP) [66]

Genomes were accessed from public databases and ANI and GBDP values calculated using online tools (<http://enve-omics.ce.gatech.edu/ani/> and <http://ggdc.dsmz.de> respectively) during December 2016. For ANI, values ≥ 95 were proposed to mimic conventional DDH results of $\geq 70\%$ [65]. For GBDP, values are proposed equivalent to conventional DDH [66]. Genome designations are GenBank reference numbers. The identity of non-type strain genomes was further verified by determining the phylogenetic relationship of their 16S rRNA gene sequences with those derived from type strains (Fig. 1 and Table S1). For *C. jejuni* subspecies, *C. coli*, *C. fetus* and *C. hyointestinalis*, MLST [155–157] was also undertaken. ANI, Average Nucleotide Identity; GBDP, Genome Blast Distance Phylogeny; SD, standard deviation; NCTC, National Collection of Type Cultures, England; CVM, Center for Veterinary Medicine, Maryland, USA; LMG, Laboratoire Microbiologie Ghent, Belgium; RM, Robert Mandrell collection, US Department of Agriculture; ATCC, American Type Culture Collection. T, type strain.

Reference taxon/strain/genome	Comparator		ANI Outputs*			GBDP outputs*		
	Taxon	Strain no.	Genome	Two way ANI	SD	No. of fragments	Formula 1	Formula 2 Formula 3
<i>Campylobacter jejuni</i> subsp. <i>jejuni</i> NCTC 11168/AL111168.1	<i>C. jejuni</i> subsp. <i>jejuni</i>	81116	NC_009839.1	97.83	1.81	6804	89.3	79.9 90.5
	<i>C. jejuni</i> subsp. <i>jejuni</i>	81-176	NC_008787.1	98.08	2.09	7029	93.9	81.6 94.3
	<i>C. jejuni</i> subsp. <i>doylei</i>	L269.97	NC_009707.1	95.97	2.3	6387	75.2	67 76.3
	<i>C. coli</i>	15-537360	NC_022660.1	85.18	6.84	4191	73.1*	27.9 58.7
	<i>C. coli</i>	CVM N29710	NC_022347.1	84.92	6.65	4142	75.5*	27.5 59.9
<i>C. coli</i> /15-537360/NC_022660.1	<i>C. coli</i>	CVM N29710	NC_022347.1	99.05	2.38	7593	97.6	89.1 97.9
<i>C. fetus</i> subsp. <i>testudinum</i> 84-112/NZ_HG004426.1	<i>C. fetus</i> subsp. <i>testudinum</i>	03-427 ^T	NC_022759.1	91.32	2.97	7293	86.6	46.6* 79.7
	<i>C. fetus</i> subsp. <i>fetus</i>	82-40	NC_008599.1	99.8	1.03	8390	94.5	97.3 96.6
	<i>C. fetus</i> subsp. <i>veneralis</i> bv. Intermedium	cfv03/293	CP006999.2	99.87	1.28	8908	98.8	97.4 99.3
	<i>C. hyointestinalis</i> subsp. <i>hyointestinalis</i>	LMG 9260	NZ_CP015575.1	82.26	6.82	2223	34.4	22.1 30.1
	<i>C. hyointestinalis</i> subsp. <i>lawsonii</i>	LMG 15993	NZ_CP015576.1	80.77	6.22	2013	28.9	21.6 26.2
<i>C. hyointestinalis</i> subsp. <i>hyointestinalis</i> LMG 9260 / NZ_CP015575.1	<i>C. hyointestinalis</i> subsp. <i>hyointestinalis</i>	LMG 15993	NZ_CP015576.1	94.65*	3.16	5862	70	57.6* 69.5
	<i>A. butzleri</i>	RM4018	NC_009850.1	78.09	4.7	1555	17.5	19.7 17.2
	<i>H. pylori</i>	Shi470	NC_010698.2	94.38*	2.59	6303	91.5	56.4* 87.3
	<i>H. pylori</i>	India7	NC_017372.1	94.85*	2.36	6580	91.4	59.4* 88.1
	<i>H. pylori</i>	SouthAfrica7	NC_017361.1	90.52*	3.16	5815	85.5	42.6* 77
<i>Aerobacter nitrofigilis</i> / DSM 7299 ^T /CP001999.1	<i>H. pylori</i>	J99	NC_000921.1	93.65*	2.44	6360	92.1	54* 87.1
	<i>H. mustelae</i>	NCTC 12198 ^T	NC_013949.1	Insufficient hits		48	12.7	28.5 13.1
	<i>H. hepaticus</i>	ATCC 51449	NC_004917.1	Insufficient hits		27	12.7	19.8 13.1
	<i>H. felis</i>	ATCC 49179 ^T	NC_014810.2	76.97	8.29	69	12.8	19.7 13.2
	<i>H. cinaedi</i>	CCUG 18818 ^T	NC_020555.1	Insufficient hits		39	12.6	28.6 13
<i>H. cinaedi</i> / CCUG 18818 ^T / NC_020555.1	<i>H. canis</i>	NCTC 12740	NZ_K1669458	83.78	8.63	285	14	26.2 14.3
	<i>H. hepaticus</i>	ATCC 51449	NC_004917.1	77.94	4.84	1126	17.7	19.8 17.4

*Denotes results that are discordant with current classifications.

with which a close phylogenetic relationship has been indicated, to determine interspecific genomic relatedness. Computational approaches described include Average Nucleotide Identity (ANI: [65]) and Genome Blast Distance Phylogeny (GBDP: [66]). Each algorithm is available online (presently <http://enve-omics.ce.gatech.edu/ani/> for ANI; and <http://ggdc.dsmz.de> for GBDP). For new species proposals, we recommend both these analyses be presented where whole-genome sequences are used as the source of the genomic data. Complete or draft sequences of appropriate comparator type strains should be used as the basis for comparison. Table 4 lists comparisons between selected complete genomes of well-characterised members of the families *Campylobacteraceae* and *Helicobacteraceae*, using the default parameters suggested online for ANI and GBDP algorithms as exemplar output. These data identify some discordance between certain comparator taxa among almost all outputs and presently-accepted classifications; Formula 3 used for GBDP analysis performed optimally on this data set (Table 4). Nonetheless, a continued validation of Formula 3 for GBDP analysis is prudent.

Description of subspecies

There is precedent for the description of genetically- defined subspecies among the *Campylobacteraceae* in particular (notably for *C. jejuni*, *C. lari*, *C. fetus* and *C. hyointestinalis*) [31, 67–69]. Most of these subspecies (apart from *C. fetus* subsp. *fetus* and *C. fetus* subsp. *venerealis*, where the definition is historically based on differing disease aetiologies) [5, 31] are characterized by a high level of infraspecific similarity and distinctive phenotype and often ecotype. This definition aligns with the concept described by Wayne *et al.* [54], whereby ‘subspecies designations can be used for genetically-close organisms that diverge in phenotype’. Subspecies should therefore exhibit DNA–DNA relatedness to the type strain of the species approximating or exceeding 70 %, with strains belonging to a given subspecies demonstrating a higher degree of genomic similarity, and clear differential characteristics in genotype, phenotype and/or ecotype among differing subspecies, as with the examples listed above.

Ecology

The natural habitat(s) of the proposed species should be detailed as much as possible, to include location(s), host species (if applicable), site of isolation, pathogenicity and clinical features (if appropriate).

Closing remarks

Minimal standards for describing new prokaryotic species aim to provide clear guidelines to the scientific community to assist in the delineation of novel taxa in a robust and unambiguous manner. This can only help the community at large who may be required to rapidly recognize emerging threats (or benefits) to public, plant, animal or environmental health. The standards described herein consider (i) previous recommendations for these taxa [8, 13, 14]; (ii) current, and indeed previous recommendations for characterisation

of prokaryotes for taxonomic purposes [54–56], and (iii) the emerging discipline of genomic taxonomy, which is in a relative state of infancy and indeed flux [70–73]. At present, the continued need for a polyphasic taxonomic approach to achieve stable and robust classifications remains critical. These standards support that need.

Funding information

The authors received no specific grant from any funding agency.

Acknowledgements

The authors would like to thank all of the members of the International Committee on Systematic Bacteriology Subcommittee on the Taxonomy of *Campylobacter* and Related Bacteria for their comments on this manuscript. This document owes much to this committee's previous publication of Minimal Standards for describing new species of the family *Campylobacteraceae* [13] and *Helicobacter* spp. [14]. Alyssa Terestre Pappa (MIT) is thanked for the updates to the Tables in the revised manuscript.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Garrity GM, Bell JA, Lilburn T. Family II. *Helicobacteraceae* fam. nov. In: Brenner DJ, Krieg NR, Staley JT, Garrity GM (editors). *Bergey's Manual of Systematic Bacteriology*, 2 ed, vol. 2, (The Proteobacteria), Part C (The Alpha-, Beta-, Delta-, and Epsilonproteobacteria). New York, USA: Springer; 2005. pp. 1168.
2. Sebald M, Véron M. Teneur en bases de L'ADN et classification des vibrios. *Annales de l'Institut Pasteur* 1963;105:897–910.
3. Goodwin CS, Armstrong JA, Chilvers T, Peters M, Collins MD *et al.* Transfer of *Campylobacter pylori* and *Campylobacter mustelae* to *Helicobacter* gen. nov. as *Helicobacter pylori* comb. nov. and *Helicobacter mustelae* comb. nov., respectively. *Int J Syst Bacteriol* 1989;39:397–405.
4. Campbell BJ, Engel AS, Porter ML, Takai K. The versatile ϵ -proteobacteria: key players in sulphidic habitats. *Nat Rev Microbiol* 2006;4:458–468.
5. On SLW. Taxonomy, phylogeny, and methods for the identification of *Campylobacter* species. In: Ketley JM and Konkel ME (editors). *Campylobacter: Molecular and Cellular Biology*. Wymondham, UK: Horizon Bioscience; 2005. pp. 13–42.
6. Vandamme P, van Doorn LJ, Al Rashid ST, Quint WG, van der Plas J *et al.* *Campylobacter hyoilei* Alderton *et al.* 1995 and *Campylobacter coli* Véron and Chatelain 1973 are subjective synonyms. *Int J Syst Bacteriol* 1997;47:1055–1060.
7. Vandamme P, Harrington CS, Jalava K, On SLW. Misidentifying helicobacters: the *Helicobacter cinaedi* example. *J Clin Microbiol* 2000;38:2261–2266.
8. Vandamme P, On SLW. Recommendations of the subcommittee on the taxonomy of *Campylobacter* and related Bacteria. *Int J Syst Evol Microbiol* 2001;51:719–721.
9. Suerbaum S, Kraft C, Dewhirst FE, Fox JG. *Helicobacter nemes-trinae* ATCC 49396^T is a strain of *Helicobacter pylori* (Marshall *et al.* 1985) Goodwin *et al.* 1989, and *Helicobacter nemes-trinae* Bronsdon *et al.* 1991 is therefore a junior heterotypic synonym of *Helicobacter pylori*. *Int J Syst Evol Microbiol* 2002;52:437–439.
10. Miller WG, On SLW. International committee on systematics of prokaryotes. subcommittee on the taxonomy of campylobacter and related bacteria: minutes of the closed meeting, 2 September 2009, Niigata, Japan. *Int J Syst Evol Microbiol* 2011;61:2559–2560.
11. On SLW. International Committee on Systematic Bacteriology Subcommittee on the taxonomy of *Campylobacter* and related Bacteria: minutes of the meeting 31st July 2002, Paris, France. *Int J Syst Evol Microbiol*;2004:291–292.

12. On SLW, Owen RJ. International committee on systematics of prokaryotes; subcommittee on the taxonomy of campylobacter and related bacteria: minutes of the meetings, 3 and 4 September 2007, Rotterdam, Holland. *Int J Syst Evol Microbiol*;2009: 197–199.
13. Ursing JB, Lior H, Owen RJ. Proposal of minimal standards for describing new species of the family *Campylobacteraceae*. *Int J Syst Bacteriol* 1994;44:842–845.
14. Dewhirst FE, Fox JG, On SLW. Recommended minimal standards for describing new species of the genus *Helicobacter*. *Int J Syst Evol Microbiol* 2000;50:2231–2237.
15. Zhang Y, Sievert SM. Pan-genome analyses identify lineage- and niche-specific markers of evolution and adaptation in *Epsilonproteobacteria*. *Front Microbiol* 2014;5:110.
16. Sasi Jyothsna TS, Rahul K, Ramaprasad EV, Sasikala C, Ramana C. *Arcobacter anaerophilus* sp. nov., isolated from an estuarine sediment and emended description of the genus *Arcobacter*. *Int J Syst Evol Microbiol* 2013;63:4619–4625.
17. On SLW, Holmes B, Sackin MJ. A probability matrix for the identification of campylobacters, helicobacters and allied taxa. *J Appl Bacteriol* 1996;81:425–432.
18. Hugh R, Leifson E. The taxonomic significance of fermentative versus oxidative metabolism of carbohydrates by various gram negative bacteria. *J Bacteriol* 1953;66:24–26.
19. Dewhirst FE, Seymour C, Fraser GJ, Paster BJ, Fox JG. Phylogeny of *Helicobacter* isolates from bird and swine feces and description of *Helicobacter pametensis* sp. nov. *Int J Syst Bacteriol* 1994;44:553–560.
20. Gill J, Haydon TG, Rawdon TG, Mcfadden AM, Ha HJ et al. *Helicobacter bilis* and *Helicobacter trogonum*: infectious causes of abortion in sheep. *J Vet Diagn Invest* 2016;28:225–234.
21. Paster BJ, Dewhirst FE. Phylogeny of Campylobacters, Wolinellas, *Bacteroides gracilis*, and *Bacteroides ureolyticus* by 16S ribosomal ribonucleic acid sequencing. *Int J Syst Bacteriol* 1988;38: 56–62.
22. Miller WG, Yee E, Jolley KA, Chapman MH. Use of an improved *atpA* amplification and sequencing method to identify members of the *Campylobacteraceae* and *Helicobacteraceae*. *Lett Appl Microbiol* 2014;58:582–590.
23. Korczak BM, Stieber R, Emler S, Burnens AP, Frey J et al. Genetic relatedness within the genus *Campylobacter* inferred from *rpoB* sequences. *Int J Syst Evol Microbiol* 2006;56:937–945.
24. Kärenlampi RI, Tolvanen TP, Hänninen ML. Phylogenetic analysis and PCR-restriction fragment length polymorphism identification of *Campylobacter* species based on partial *groEL* gene sequences. *J Clin Microbiol* 2004;42:5731–5738.
25. Megraud F, Bonnet F, Garnier M, Lamouliatte H. Characterization of “*Campylobacter pyloridis*” by culture, enzymatic profile, and protein content. *J Clin Microbiol* 1985;22:1007–1010.
26. On SLW, Holmes B. Effect of inoculum size on the phenotypic characterisation of *Campylobacter* spp. *J Clin Micro* 1991a;29: 923–926.
27. On SLW, Holmes B. Reproducibility of tolerance tests that are useful in the identification of campylobacteria. *J Clin Microbiol* 1991;29:1785–1788.
28. On SLW, Holmes B. Assessment of enzyme detection tests useful in identification of campylobacteria. *J Clin Microbiol* 1992;30: 746–749.
29. On SLW, Holmes B. Classification and identification of campylobacters, helicobacters and allied taxa by numerical analysis of phenotypic characters. *System Appl Microbiol* 1995;18:374–390.
30. Alispahic M, Hummel K, Jandreski-Cvetkovic D, Nöbauer K, Razzazi-Fazeli E et al. Species-specific identification and differentiation of *Arcobacter*, *helicobacter* and *Campylobacter* by full-spectral matrix-associated laser desorption/ionization time of flight mass spectrometry analysis. *J Med Microbiol* 2010;59: 295–301.
31. Fitzgerald C, Tu ZC, Patrick M, Stiles T, Lawson AJ et al. *Campylobacter fetus* subsp. *testudinum* subsp. nov., isolated from humans and reptiles. *Int J Syst Evol Microbiol* 2014;64:2944–2948.
32. Levican A, Rubio-Arcos S, Martinez-Murcia A, Collado L, Figueras MJ. *Arcobacter ebronensis* sp. nov. and *Arcobacter aquimarinus* sp. nov., two new species isolated from marine environment. *Syst Appl Microbiol* 2015;38:30–35.
33. Mandrell RE, Harden LA, Bates A, Miller WG, Haddon WF et al. Speciation of *Campylobacter coli*, *C. jejuni*, *C. helveticus*, *C. lari*, *C. sputorum*, and *C. upsaliensis* by matrix-assisted laser desorption ionization-time of flight mass spectrometry. *Appl Environ Microbiol* 2005;71:6292–6307.
34. Bessède E, Solecki O, Sifré E, Labadi L, Mégraud F. Identification of *Campylobacter* species and related organisms by matrix assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrometry. *Clin Microbiol Infect* 2011;17:1735–1739.
35. Murray RG, Schleifer KH. Taxonomic notes: a proposal for recording the properties of putative taxa of procaryotes. *Int J Syst Bacteriol* 1994;44:174–176.
36. Murray RG, Stackebrandt E. Taxonomic note: implementation of the provisional status candidatus for incompletely described procaryotes. *Int J Syst Bacteriol* 1995;45:186–187.
37. Lawson AJ, Linton D, Stanley J. 16s rRNA gene sequences of ‘*Candidatus Campylobacter hominis*’, a novel uncultivated species, are found in the gastrointestinal tract of healthy humans. *Microbiology* 1998;144:2063–2071.
38. Cook GT. A plate test for Nitrate Reduction. *J Clin Pathol* 1950;3: 359–362.
39. Owen RJ, Martin SR, Borman P. Rapid urea hydrolysis by gastric campylobacters. *Lancet* 1985;1:111.
40. Itoh T, Yanagawa Y, Shingaki M, Takahashi M, Kai A et al. Isolation of *Campylobacter pyloridis* from human gastric mucosa and characterization of the isolates. *Microbiol Immunol* 1987;31:603–614.
41. Skirrow MB, Benjamin J. Differentiation of enteropathogenic campylobacter. *J Clin Pathol* 1980;33:1122.
42. van den Bulck K, Decostere A, Baele M, Vandamme P, Mast J et al. *Helicobacter cynogastricus* sp. nov., isolated from the canine gastric mucosa. *Int J Syst Evol Microbiol* 2006;56:1559–1564.
43. Hawrylik SJ, Wasilko DJ, Haskell SL, Gootz TD, Lee SE. Bisulfite or sulfite inhibits growth of *Helicobacter pylori*. *J Clin Microbiol* 1994;32:790–792.
44. Fox JG, Yan LL, Dewhirst FE, Paster BJ, Shames B et al. *Helicobacter bilis* sp. nov., a novel *Helicobacter* species isolated from bile, livers, and intestines of aged, inbred mice. *J Clin Microbiol* 1995;33:445–454.
45. Harrington CS, On SLW. Extensive 16S rRNA gene sequence diversity in *Campylobacter hyointestinalis* strains: taxonomic and applied implications. *Int J Syst Bacteriol* 1999;49:1171–1175.
46. Hänninen ML, Kärenlampi RI, Koort JM, Mikkonen T, Björkroth KJ. Extension of the species *Helicobacter bilis* to include the reference strains of *Helicobacter* sp. flexispira taxa 2, 3 and 8 and finnish canine and feline flexispira strains. *Int J Syst Evol Microbiol* 2005;55:891–898.
47. Hänninen ML, Utriainen M, Happonen I, Dewhirst FE. *Helicobacter* sp. flexispira 16S rDNA taxa 1, 4 and 5 and finnish porcine *Helicobacter* isolates are members of the species *Helicobacter trogonum* (taxon 6). *Int J Syst Evol Microbiol* 2003; 53:425–433.
48. Glaeser SP, Kämpfer P. Multilocus sequence analysis (MLSA) in prokaryotic taxonomy. *Syst Appl Microbiol* 2015;38:237–245.
49. Jolley KA, Bliss CM, Bennett JS, Bratcher HB, Brehony C et al. Ribosomal multilocus sequence typing: universal characterization of Bacteria from domain to strain. *Microbiology* 2012;158: 1005–1015.

50. Levican A, Collado L, Figueras MJ. *Arcobacter cloacae* sp. nov. and *Arcobacter suis* sp. nov., two new species isolated from food and sewage. *Syst Appl Microbiol* 2013;36:22–27.
51. Ankenbrand MJ, Keller A, Chain F. bcgTree: automatized phylogenetic tree building from bacterial core genomes. *Genome* 2016;59:783–791.
52. Zhang Y, Qiu S. Phylogenomic analysis of the genus *Ralstonia* based on 686 single-copy genes. *Antonie van Leeuwenhoek* 2016;109:71–82.
53. de Ley J, Cattoir H, Reynaerts A. The quantitative measurement of DNA hybridization from renaturation rates. *Eur J Biochem* 1970;12:133–142.
54. Wayne LG, Moore WEC, Stackebrandt E, Kandler O, Colwell RR et al. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Evol Microbiol* 1987;37:463–464.
55. Stackebrandt E, Frederiksen W, Garrity GM, Grimont PA, Kämpfer P et al. Report of the ad hoc committee for the re-evaluation of the species definition in bacteriology. *Int J Syst Evol Microbiol* 2002;52:1043–1047.
56. Tindall BJ, Rosselló-Móra R, Busse HJ, Ludwig W, Kämpfer P. Notes on the characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol* 2010;60:249–266.
57. Fox JG, Taylor NS, Edmonds P, Brenner DJ. *Campylobacter pylori* subsp. *mustelae* subsp. nov. isolated from the gastric mucosa of ferrets (*Mustela putorius furo*), and an emended description of *Campylobacter pylori*. *Int J Syst Bacteriol* 1988;38:367–370.
58. Hänninen M-L, Happonen I, Saari S, Jalava K. Culture and characteristics of *Helicobacter bizzozeronii*, a new canine gastric *Helicobacter* sp. *Int J Syst Bacteriol* 46, 160–166. Erratum in: *Int J Syst Bacteriol* 1996;46:839.
59. Owen RJ, Pitcher D. Current methods for estimating DNA base composition and levels of DNA–DNA hybridization. In: Goodfellow M and Minnikin DE (editors). *Chemical Methods in Bacterial Systematics*. London, UK: Academic Press; 1985. pp. 67–93.
60. Stanley J, Burnens AP, Linton D, On SL, Costas M et al. *Campylobacter helveticus* sp. nov., a new thermophilic species from domestic animals: characterization, and cloning of a species-specific DNA probe. *J Gen Microbiol* 1992;138:2293–2303.
61. Ezaki T, Takeuchi N, Liu SL, Kai A, Yamamoto H et al. Small-scale DNA preparation for rapid genetic identification of *Campylobacter* species without radioisotope. *Microbiol Immunol* 1988;32:141–150.
62. Fox JG, Chilvers T, Goodwin CS, Taylor NS, Edmonds P et al. *Campylobacter mustelae*, a new species resulting from the elevation of *Campylobacter pylori* subsp. *mustelae* to Species Status. *Int J Syst Bacteriol* 1989;39:301–303.
63. On SLW, Harrington CS. Identification of taxonomic and epidemiological relationships among *Campylobacter* species by numerical analysis of AFLP profiles. *FEMS Microbiol Lett* 2000;193:161–169.
64. On SLW, Harrington CS, Atabay HI. Differentiation of *Arcobacter* species by numerical analysis of AFLP profiles and description of a novel *Arcobacter* from pig abortions and Turkey faeces. *J Appl Microbiol* 2003;95:1096–1105.
65. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P et al. DNA–DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 2007;57:81–91.
66. Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 2013;14:60.
67. Steele TW, Owen RJ. NOTES: *Campylobacter jejuni* subsp. *doylei* subsp. nov., a subspecies of nitrate-negative campylobacters isolated from human clinical specimens. *Int J Syst Bacteriol* 1988;38:316–318.
68. Debruyne L, On SLW, de Brandt E, Vandamme P. Novel *Campylobacter lari*-like bacteria from humans and molluscs: description of *Campylobacter peloridis* sp. nov., *Campylobacter lari* subsp. *concheus* subsp. nov. and *Campylobacter lari* subsp. *lari* subsp. nov. *Int J Syst Evol Microbiol* 2009;59:1126–1132.
69. On SLW, Bloch B, Holmes B, Hoste B, Vandamme P. *Campylobacter hyointestinalis* subsp. *lawsonii* subsp. nov., isolated from the porcine stomach, and an emended description of *Campylobacter hyointestinalis*. *Int J Syst Bacteriol* 1995;45:767–774.
70. Gevers D, Cohan FM, Lawrence JG, Spratt BG, Coenye T et al. Opinion: re-evaluating prokaryotic species. *Nat Rev Microbiol* 2005;3:733–739.
71. Thompson CC, Chimetto L, Edwards RA, Swings J, Stackebrandt E et al. Microbial genomic taxonomy. *BMC Genomics* 2013;14:913.
72. Chun J, Rainey FA. Integrating genomics into the taxonomy and systematics of the bacteria and archaea. *Int J Syst Evol Microbiol* 2014;64:316–324.
73. Rosselló-Móra R, Amann R. Past and future species definitions for bacteria and archaea. *Syst Appl Microbiol* 2015;38:209–216.
74. Collado L, Cleenwerck I, van Trappen S, de Vos P, Figueras MJ. *Arcobacter mytili* sp. nov., an indoxyl acetate-hydrolysis-negative bacterium isolated from mussels. *Int J Syst Evol Microbiol* 2009;59:1391–1396.
75. de Smet S, Vandamme P, De Zutter L, On SLW, Doudiah L et al. *Arcobacter trophiarum* sp. nov., isolated from fattening pigs. *Int J Syst Evol Microbiol* 2011;61:356–361.
76. Donachie SP, Bowman JP, On SL, Alam M. *Arcobacter halophilus* sp. nov., the first obligate halophile in the genus *Arcobacter*. *Int J Syst Evol Microbiol* 2005;55:1271–1277.
77. Figueras MJ, Levican A, Collado L, Inza MI, Yustes C. *Arcobacter ellisii* sp. nov., isolated from mussels. *Syst Appl Microbiol* 2011;34:414–418.
78. Figueras MJ, Collado L, Levican A, Perez J, Solsona MJ et al. *Arcobacter molluscorum* sp. nov., a new species isolated from shellfish. *Syst Appl Microbiol* 2011;34:105–109.
79. Houf K, On SLW, Coenye T, Debruyne L, de Smet S et al. *Arcobacter thereius* sp. nov., isolated from pigs and ducks. *Int J Syst Evol Microbiol* 2009;59:2599–2604.
80. Houf K, On SLW, Coenye T, Mast J, van Hoof J et al. *Arcobacter cibarius* sp. nov., isolated from broiler carcasses. *Int J Syst Evol Microbiol* 2005;55:713–717.
81. Kiehlbauch JA, Brenner DJ, Nicholson MA, Baker CN, Patton CM et al. *Campylobacter butzleri* sp. nov. isolated from humans and animals with diarrheal illness. *J Clin Microbiol* 1991;29:376–385.
82. Kim HM, Hwang CY, Cho BC. *Arcobacter marinus* sp. nov. *Int J Syst Evol Microbiol* 2010;60:531–536.
83. Levican A, Collado L, Aguilar C, Yustes C, Diéguez AL et al. *Arcobacter bivalviorum* sp. nov. and *Arcobacter venerupis* sp. nov., new species isolated from shellfish. *Syst Appl Microbiol* 2012;35:133–138.
84. Neill SD, Campbell JN, O'Brien JJ, Weatherup STC, Ellis WA. Taxonomic position of *Campylobacter cryaerophila* sp. nov. *Int J Syst Bacteriol* 1985;35:342–356.
85. Mcclung CR, Patriquin DG, Davis RE. *Campylobacter nitrofigilis* sp. nov., a nitrogen-fixing bacterium associated with roots of *Spartina alterniflora* Loisel. *Int J Syst Bacteriol* 1983;33:605–612.
86. Vandamme P, Vancanneyt M, Pot B, Mels L, Hoste B et al. Polyphasic taxonomic study of the emended genus *Arcobacter* with *Arcobacter butzleri* comb. nov. and *Arcobacter skirrowii* sp. nov., an aerotolerant bacterium isolated from veterinary specimens. *Int J Syst Bacteriol* 1992;42:344–356.
87. Vandamme P, Falsen E, Rossau R, Hoste B, Segers P et al. Revision of *Campylobacter*, *helicobacter*, and *Wolinella* taxonomy: emendation of generic descriptions and proposal of *Arcobacter* gen. nov. *Int J Syst Bacteriol* 1991;41:88–103.

88. Whiteduck-Léveillé K, Whiteduck-Léveillé J, Cloutier M, Tambong JT, Xu R et al. *Arcobacter lanthieri* sp. nov., isolated from pig and dairy cattle manure. *Int J Syst Evol Microbiol* 2015; 65:2709–2716.
89. Benjamin J, Leaper S, Owen RJ, Skirrow MB. Description of *Campylobacter laridis*, a new species comprising the nalidixic acid resistant thermophilic *Campylobacter* (NARTC) group. *Curr Microbiol* 1983;8:231–238.
90. Cáceres A, Muñoz I, Iraola G, Díaz-Viraqué F, Collado L. *Campylobacter ornithocola* sp. nov., a novel member of the *Campylobacter lari* group isolated from wild bird faecal samples. *Int J Syst Evol Microbiol* 2017;67:1643–1649.
91. Debruyne L, Broman T, Bergström S, Olsen B, On SLW et al. *Campylobacter subantarcticus* sp. nov., isolated from birds in the sub-Antarctic region. *Int J Syst Evol Microbiol* 2010;60:815–819.
92. Debruyne L, Broman T, Bergström S, Olsen B, On SL et al. *Campylobacter volucris* sp. nov., isolated from black-headed gulls (*Larus ridibundus*). *Int J Syst Evol Microbiol* 2010;60:1870–1875.
93. Doyle LP. The etiology of swine dysentery. *Am J Vet Res* 1948;9: 50–51.
94. Florent A. Les deux vibrioses génitales: la vibriose due à *V. fetus venerealis* et la vibriose d'origine intestinale due à *V. fetus intestinalis*. *Meded Veeartsenijsh Rijksuniv Gent* 1959;3:1–60.
95. Foster G, Holmes B, Steigerwalt AG, Lawson PA, Thorne P et al. *Campylobacter insulaenigrae* sp. nov., isolated from marine mammals. *Int J Syst Evol Microbiol* 2004; 54:2369–2373. Erratum in: *Int J Syst Evol Microbiol* 2005;55:981.
96. Gebhart CJ, Edmonds P, Ward GE, Kurtz HJ, Brenner DJ. "*Campylobacter hyointestinalis*" sp. nov.: a new species of *Campylobacter* found in the intestines of pigs and other animals. *J Clin Microbiol* 1985;21:715–720.
97. Gilbert MJ, Kik M, Miller WG, Duim B, Wagenaar JA. *Campylobacter iguaniorum* sp. nov., isolated from reptiles. *Int J Syst Evol Microbiol* 2015;65:975–982.
98. Gilbert MJ, Miller WG, Leger JS, Chapman MH, Timmerman AJ et al. *Campylobacter pinnipediorum* sp. nov., isolated from pinnipeds, comprising *Campylobacter pinnipediorum* subsp. *pinnipediorum* subsp. nov. and *Campylobacter pinnipediorum* subsp. *caledonicus* subsp. nov. *Int J Syst Evol Microbiol* 2017;67:1961–1968.
99. Inglis GD, Hoar BM, Whiteside DP, Morck DW. *Campylobacter canadensis* sp. nov., from captive whooping cranes in Canada. *Int J Syst Evol Microbiol* 2007;57:2636–2644.
100. Jackson FL, Goodman YE. *Bacteroides ureolyticus*, a new species to accommodate strains previously identified as "*Bacteroides corrodens*, Anaerobic". *Int J Syst Bacteriol* 1978;28:197–200.
101. Jones FS, Orcutt M, Little RB. Vibrios (*Vibrio jejuni*, n. sp.) associated with intestinal disorders of cows and calves. *J Exp Med* 1931;53:853–863.
102. Koziel M, O'Doherty P, Vandamme P, Corcoran GD, Sleator RD et al. *Campylobacter corcagiensis* sp. nov., isolated from faeces of captive lion-tailed macaques (*Macaca silenus*). *Int J Syst Evol Microbiol* 2014;64:2878–2883.
103. Lawson AJ, On SLW, Logan JM, Stanley J. *Campylobacter hominis* sp. nov., from the human gastrointestinal tract. *Int J Syst Evol Microbiol* 2001;51:651–660.
104. Lawson GH, Rowland AC. Intestinal adenomatosis in the pig: a bacteriological study. *Res Vet Sci* 1974;17:331–336.
105. Lawson GHK, Leaver JL, Pettigrew GW, Rowland AC. Some features of *Campylobacter sputorum* subsp. *mucosalis* subsp. nov., nom. rev. and their taxonomic significance. *Int J Syst Bacteriol* 1981;31:385–391.
106. Logan JM, Burnens A, Linton D, Lawson AJ, Stanley J. *Campylobacter lanienae* sp. nov., a new species isolated from workers in an abattoir. *Int J Syst Evol Microbiol* 2000;50:865–872.
107. On SLW, Atabay HI, Corry JE, Harrington CS, Vandamme P. Emended description of *Campylobacter sputorum* and revision of its infrasubspecific (biovar) divisions, including *C. sputorum* biovar *paraureolyticus*, a urease-producing variant from cattle and humans. *Int J Syst Bacteriol* 1998;48:195–206.
108. Prévot AR. Études de systématique bactérienne. V. Essai de classification des vibrions anaérobies. *Ann Inst Pasteur* 1940;64: 117–125.
109. Roop RM, Smibert RM, Johnson JL, Krieg NR. *Campylobacter mucosalis* (Lawson, Leaver, Pettigrew, and Rowland 1981) comb. nov.: emended description. *Int J Syst Bacteriol* 1985;35: 189–192.
110. Rossi M, Debruyne L, Zanoni RG, Manfreda G, Revez J et al. *Campylobacter avium* sp. nov., a hippurate-positive species isolated from poultry. *Int J Syst Evol Microbiol* 2009;59:2364–2369.
111. Sandstedt K, Ursing J. Description of *Campylobacter upsaliensis* sp. nov. previously known as the CNW Group. *Syst Appl Microbiol* 1991;14:39–45.
112. Smith T, Taylor MS. Some morphological and biological characters of the spirilla (*Vibrio fetus*, N. sp.) associated with disease of the fetal membranes in cattle. *J Exp Med* 1919;30:299–311.
113. Vandamme P, Debruyne L, de Brandt E, Falsen E. Reclassification of *Bacteroides ureolyticus* as *Campylobacter ureolyticus* comb. nov., and emended description of the genus *Campylobacter*. *Int J Syst Evol Microbiol* 2010;60:2016–2022.
114. Vandamme P, Daneshvar MI, Dewhirst FE, Paster BJ, Kersters K et al. Chemotaxonomic analyses of *Bacteroides gracilis* and *Bacteroides ureolyticus* and reclassification of *B. gracilis* as *Campylobacter gracilis* comb. nov. *Int J Syst Bacteriol* 1995;45:145–152.
115. Veron M, Chatelain R. Taxonomic study of the genus *Campylobacter* sebal and Veron and designation of the neotype strain for the type species, *Campylobacter fetus* (Smith and Taylor) Sebal and Veron. *Int J Syst Bacteriol* 1973;23:122–134.
116. van TT, Elshagmani E, Gor MC, Scott PC, Moore RJ. *Campylobacter hepaticus* sp. nov., isolated from chickens with spotty liver disease. *Int J Syst Evol Microbiol* 2016;66:4518–4524.
117. von Graevenitz A. Revised nomenclature of *Campylobacter lari-dis*, *Enterobacter intermedium*, and "*Flavobacterium branchiophila*". *Int J Syst Bacteriol* 1990;40:211.
118. Tanner ACR, Badger S, Lai C-H, Listgarten MA, Visconti RA et al. *Wolinella* gen. nov., *Wolinella succinogenes* (*Vibrio succinogenes* Wolin et al.) comb. nov., and description of *Bacteroides gracilis* sp. nov., *Wolinella recta* sp. nov., *Campylobacter concisus* sp. nov., and *Eikenella corrodens* from humans with periodontal disease. *Int J Syst Bacteriol* 1981;31:432–445.
119. Tanner ACR, Listgarten MA, Ebersole JL. *Wolinella curva* sp. nov.: "*Vibrio succinogenes*" of human origin. *Int J Syst Bacteriol* 1984;34:275–282.
120. Zanoni RG, Debruyne L, Rossi M, Revez J, Vandamme P. *Campylobacter cuniculorum* sp. nov., from rabbits. *Int J Syst Evol Microbiol* 2009;59:1666–1671.
121. Baele M, Decostere A, Vandamme P, Ceelen L, Hellemans A et al. Isolation and characterization of *Helicobacter suis* sp. nov. from pig stomachs. *Int J Syst Evol Microbiol* 2008;58:1350–1358.
122. Baele M, Decostere A, Vandamme P, van den Bulck K, Gruntar I et al. *Helicobacter baculiformis* sp. nov., isolated from feline stomach mucosa. *Int J Syst Evol Microbiol* 2008;58:357–364.
123. Eaton KA, Dewhirst FE, Radin MJ, Fox JG, Paster BJ et al. *Helicobacter acinonyx* sp. nov., isolated from cheetahs with gastritis. *Int J Syst Bacteriol* 1993;43:99–106.
124. Fox JG, Boutin SR, Handt LK, Taylor NS, Xu S et al. Isolation and characterization of a novel *Helicobacter* species, "*Helicobacter macacae*," from rhesus monkeys with and without chronic idiopathic colitis. *J Clin Microbiol* 2007;45:4061–4063.
125. Fox JG, Chien CC, Dewhirst FE, Paster BJ, Shen Z et al. *Helicobacter canadensis* sp. nov., isolated from humans with diarrhea as an example of an emerging pathogen. *J Clin Microbiol* 2000; 38:2546–2549.

126. Fox JG, Dewhirst FE, Tully JG, Paster BJ, Yan L et al. *Helicobacter hepaticus* sp. nov., a microaerophilic bacterium isolated from livers and intestinal mucosal scrapings from mice. *J Clin Microbiol* 1994;32:1238–1245.
127. Fox JG, Shen Z, Xu S, Feng Y, Dangler CA et al. *Helicobacter marmotae* sp. nov. isolated from livers of woodchucks and intestines of cats. *J Clin Microbiol* 2002;40:2513–2519.
128. Fox JG, Taylor NS, Howe S, Tidd M, Xu S et al. *Helicobacter anseris* sp. nov. and *Helicobacter brantae* sp. nov., isolated from feces of resident Canada geese in the greater Boston area. *Appl Environ Microbiol* 2006;72:4633–4637.
129. Franklin CL, Beckwith CS, Livingston RS, Riley LK, Gibson SV et al. Isolation of a novel *Helicobacter* species, *Helicobacter cholecystus* sp. nov., from the gallbladders of Syrian hamsters with cholangiofibrosis and centrilobular pancreatitis. *J Clin Microbiol* 1996;34:2952–2958.
130. Franklin CL, Gorelick PL, Riley LK, Dewhirst FE, Livingston RS et al. *Helicobacter typhlonius* sp. nov., a novel murine urease-negative *Helicobacter* species. *J Clin Microbiol* 2001;39:3920–3926.
131. Harper CG, Feng Y, Xu S, Taylor NS, Kinsel M et al. *Helicobacter cetorum* sp. nov., a urease-positive *Helicobacter* species isolated from dolphins and whales. *J Clin Microbiol* 2002;40:4536–4543.
132. Hu S, Jin D, Lu S, Liu S, Zhang J et al. *Helicobacter himalayensis* sp. nov. isolated from gastric mucosa of *Marmota himalayana*. *Int J Syst Evol Microbiol* 2015;65:1719–1725.
133. Jalava K, Kaartinen M, Utriainen M, Happonen I, Hänninen ML. *Helicobacter salomonis* sp. nov., a canine gastric *Helicobacter* sp. related to *Helicobacter felis* and *Helicobacter bizzozeronii*. *Int J Syst Bacteriol* 1997;47:975–982.
134. Jalava K, On SLW, Vandamme PA, Happonen I, Sukura A et al. Isolation and identification of *Helicobacter* spp. from canine and feline gastric mucosa. *Appl Environ Microbiol* 1998;64:3998–4006.
135. Lee A, Phillips MW, O'Rourke JL, Paster BJ, Dewhirst FE et al. *Helicobacter muridarum* sp. nov., a microaerophilic helical bacterium with a novel ultrastructure isolated from the intestinal mucosa of rodents. *Int J Syst Bacteriol* 1992;42:27–36.
136. Marshall BJ, Royce H, Annear DI, Goodwin CS, Pearman JW et al. Original isolation of *Campylobacter pyloridis* from human gastric mucosa. *Microbios Letters* 1984;25:83–88.
137. Marshall BJ, Goodwin CS. Notes: revised nomenclature of *Campylobacter pyloridis*. *Int J Syst Bacteriol* 1987;37:68.
138. Mendes EN, Queiroz DM, Dewhirst FE, Paster BJ, Moura SB et al. *Helicobacter trogonum* sp. nov., isolated from the rat intestine. *Int J Syst Bacteriol* 1996;46:916–921.
139. Moyaert H, Decostere A, Vandamme P, Debruyne L, Mast J et al. *Helicobacter equorum* sp. nov., a urease-negative *Helicobacter* species isolated from horse faeces. *Int J Syst Evol Microbiol* 2007;57:213–218.
140. Paster BJ, Lee A, Fox JG, Dewhirst FE, Tordoff LA et al. Phylogeny of *Helicobacter felis* sp. nov., *Helicobacter mustelae*, and related bacteria. *Int J Syst Bacteriol* 1991;41:31–38.
141. Patterson MM, Schrenzel MD, Feng Y, Xu S, Dewhirst FE et al. *Helicobacter aurati* sp. nov., a urease-positive *Helicobacter* species cultured from gastrointestinal tissues of syrian hamsters. *J Clin Microbiol* 2000;38:3722–3728.
142. Robertson BR, O'Rourke JL, Vandamme P, On SLW, Lee A. *Helicobacter ganmani* sp. nov., a urease-negative anaerobe isolated from the intestines of laboratory mice. *Int J Syst Evol Microbiol* 2001;51:1881–1889.
143. Shen Z, Fox JG, Dewhirst FE, Paster BJ, Foltz CJ et al. *Helicobacter rodentium* sp. nov., a urease-negative *Helicobacter* species isolated from laboratory mice. *Int J Syst Bacteriol* 1997;47:627–634.
144. Shen Z, Xu S, Dewhirst FE, Paster BJ, Pena JA et al. A novel enterohepatic *Helicobacter* species '*Helicobacter mastomyrinus*' isolated from the liver and intestine of rodents. *Helicobacter* 2005;10:59–70.
145. Shen Z, Feng Y, Sheh A, Everitt J, Bertram F et al. Isolation and characterization of a novel *Helicobacter* species, *Helicobacter jaa-chi* sp. nov., from common marmosets (*Callithrix jacchus*). *J Med Microbiol* 2015;64:1063–1073.
146. Shen Z, Feng Y, Muthupalani S, Sheh A, Cheaney LE et al. Novel *Helicobacter* species *H.japonicum* isolated from laboratory mice from Japan induces typhlocolitis and lower bowel carcinoma in C57BL/129 IL10^{-/-} mice. *Carcinogenesis* 2016;37:1190–1198.
147. Shen Z, Mannion A, Whary MT, Muthupalani S, Sheh A et al. *Helicobacter saguini*, a Novel *Helicobacter* isolated from cotton-top tamarins with ulcerative colitis, has proinflammatory properties and induces typhlocolitis and dysplasia in Gnotobiotic IL-10^{-/-} Mice. *Infect Immun* 2016;84:2307–2316.
148. Simmons JH, Riley LK, Besch-Williford CL, Franklin CL. *Helicobacter mesocricetorum* sp. nov., a novel *Helicobacter* isolated from the feces of Syrian hamsters. *J Clin Microbiol* 2000;38:1811–1817.
149. Smet A, Flahou B, D'Herde K, Vandamme P, Cleenwerck I et al. *Helicobacter heilmannii* sp. nov., isolated from feline gastric mucosa. *Int J Syst Evol Microbiol* 2012; 62:299–306. Erratum in: *Int J Syst Evol Microbiol* 2012;62:1016.
150. Stanley J, Linton D, Burnens AP, Dewhirst FE, On SL et al. *Helicobacter pullorum* sp. nov.-genotype and phenotype of a new species isolated from poultry and from human patients with gastroenteritis. *Microbiology* 1994;140:3441–3449.
151. Stanley J, Linton D, Burnens AP, Dewhirst FE, Owen RJ et al. *Helicobacter canis* sp. nov., a new species from dogs: an integrated study of phenotype and genotype. *J Gen Microbiol* 1993; 139:2495–2504.
152. Totten PA, Fennell CL, Tenover FC, Wezenberg JM, Perine PL et al. *Campylobacter cinaedi* (sp. nov.) and *Campylobacter fennelliae* (sp. nov.): two new *Campylobacter* species associated with enteric disease in homosexual men. *J Infect Dis* 1985;151:131–139.
153. Truper HG, De'clari L. Taxonomic note: necessary correction of specific epithets formed as substantives (Nouns) "in Apposition". *Int J Syst Bacteriol* 1997;47:908–909.
154. Wolin MJ, Wolin EA, Jacobs NJ. Cytochrome-producing anaerobic vibrio, *Vibrio succinogenes*, sp. nov. *J Bacteriol* 1961;81:911–917.
155. Miller WG, Chapman MH, Yee E, On SLW, McNulty DK et al. Multilocus sequence typing methods for the emerging *Campylobacter* species *C. hyointestinalis*, *C. lanienae*, *C. sputorum*, *C. concisus*, and *C. curvus*. *Front Cell Infect Microbiol* 2012;2:45.
156. Miller WG, On SLW, Wang G, Fontanoz S, Lastovica AJ et al. Extended multilocus sequence typing system for *Campylobacter coli*, *C. lari*, *C. upsaliensis*, and *C. helveticus*. *J Clin Microbiol* 2005; 43:2315–2329.
157. van Bergen MA, Dingle KE, Maiden MC, Newell DG, van der Graaf-van Bloois L et al. Clonal nature of *Campylobacter fetus* as defined by multilocus sequence typing. *J Clin Microbiol* 2005;43: 5888–5898.